

# **QUICK REFERENCE**

# TRANSTHORACIC ECHO SCREENING



# See Important Safety Information referenced within.

©2020 Abbott. All rights reserved. MAT-2001330 v1.0 Item approved for U.S. use only.



The following transthoracic echo (TTE) views represent key considerations for MitraClip<sup>™</sup> G4 Therapy. Adherence to this systematic protocol is recommended to ensure efficient analysis of the mitral valve and to assess anatomic eligibility for the MitraClip<sup>™</sup> G4 Procedure.

# **GENERAL COMMENTS**

- Digital archived images should include three (3) or more cardiac cycles—unless patient has atrial fibrillation, then five (5) cardiac cycles are recommended
- Ensure color Doppler Nyquist limits range from 0.5–0.7 m/sec–unless specified for PISA
- •Adjust gain and depth to enhance and maximize the image for measurements
- Perform all spectral Doppler and M-mode recordings at a sweep speed of 100 mm/sec
- Use of color compare setting is strongly recommended
- Ensure that peak spectral velocities are fully visible on screen
- Confirm that EKG signal is clearly visible on all frames
- •All calibration lines should be clearly visible
- •Use of a customized echocardiography bed is strongly recommended
- •Use 3D images to supplement and confirm initial diagnosis
- Ensure that all cardiac structures are analyzed per institution guidelines

# See Important Safety Information referenced within.

## TTE ASSESSMENT CHECKLIST

# 1 COLOR FLOW JET

# □ None

- □ Mild
- □ Moderate □ Moderate-to-severe

□ Severe

## PULMONARY **VEIN FLOW**

□ Normal pulmonary vein flow

□ Codominant pulmonary vein flow

Diastolic dominant pulmonary vein flow

Systolic pulmonary vein flow reversal

**VENA CONTRACTA** WIDTH (cm)

- REGURGITANT VOLUME (ml/beat)
- **5** REGURGITANT FRACTION (%)
- REGURGITANT ORIFICE AREA (cm<sup>2</sup>)

# PARASTERNAL LONG AXIS VIEW



### 7 MITRAL VALVE ORIFICE AREA (cm<sup>2</sup>)

LV EJECTION **FRACTION (%)** 

> LV END SYSTOLIC **DIMENSION (LVIDS)** (cm)

## **PRESENCE OF** MITRAL ANNULAR **CALCIFICATIONS**

□ None □ Mild/moderate Severe

### **ORIGIN OF PRIMARY REGURGITANT JET**

**PRESENCE OF A** SECOND CLINICALLY **SIGNIFICANT JET** 

**MR ETIOLOGY** 

□ Secondary □ Primary □ Mixed

IN THIS VIEW, ASSESS:

• calcification in mitral

vena contracta width

• A2/P2 pathology

valve area (if any/severity)

• LV size and function

• LA size

• MR etiology

## IN THIS VIEW, ASSESS: • for ASDs. VSDs. and

PARASTERNAL SHORT AXIS VIEW: AORTIC VALVE LEVEL



**APICAL 4-CHAMBER VIEW** 

APICAL 5-CHAMBER VIEW

## **PARASTERNAL SHORT AXIS VIEW: MITRAL VALVE LEVEL**



## calcification in mitral valve area (if any/severity)

shunts by interrogating

the intra-atrial septum

- · jet origin with color
- Doppler applied • size of mitral valve area by planimetry

# IN THIS VIEW, ASSESS:

• LV size and function • LA size

- MR etiology
- MR severity
- · pulmonary vein flow
- calcification in mitral valve area (if any/severity)

### IN THIS VIEW, ASSESS:

- LA size
- MR etiology
- MR severity
- pulmonary vein flow
- interrogate aortic valve using standard technique







## **APICAL 2-CHAMBER VIEW**



### **APICAL 3-CHAMBER VIEW**

### IN THIS VIEW, ASSESS:

- LV size and function
- LA size
- MR etiology
- MR severity
- · pulmonary vein flow
- calcification in mitral valve area (if any/severity)
- jet origin with color Doppler applied

### IN THIS VIEW, ASSESS:

- LV size and function
- LA size
- MR etiology
- calcification in mitral valve area (if any/severity)

# SUBCOSTAL LONG AXIS VIEW



### IN THIS VIEW, ASSESS:

• color Doppler of atrial septum to interrogate presence of ASD

## SUBCOSTAL SHORT AXIS VIEW



### IN THIS VIEW, ASSESS:

• 2D of inferior vena cava collapsing (sniff test)

# IMPORTANT SAFETY INFORMATION

# **R** MITRACLIP CLIP DELIVERY SYSTEMS

### **INDICATION FOR USE**

- The MitraClip<sup>™</sup> G4 System is indicated for the percutaneous reduction of significant symptomatic mitral regurgitation (MR ≥ 3+) due to primary abnormality of the mitral apparatus [degenerative MR] in patients who have been determined to be at prohibitive risk for mitral valve surgery by a heart team, which includes a cardiac surgeon experienced in mitral valve surgery and a cardiologist experienced in mitral valve disease, and in whom existing comorbidities would not preclude the expected benefit from reduction of the mitral regurgitation.
- The MitraClip<sup>™</sup> G4 System, when used with maximally tolerated guideline-directed medical therapy (GDMT), is indicated for the treatment of symptomatic, moderate-to-severe or severe secondary (or functional) mitral regurgitation (MR; MR ≥ Grade III per American Society of Echocardiography criteria) in patients with a left ventricular ejection fraction (LVEF) ≥ 20% and ≤ 50%, and a left ventricular end systolic dimension (LVESD) ≤ 70 mm whose symptoms and MR severity persist despite maximally tolerated GDMT as determined by a multidisciplinary heart team experienced in the evaluation and treatment of heart failure and mitral valve disease.

### CONTRAINDICATIONS

The MitraClip  $^{\rm TM}$  G4 System is contraindicated in patients with the following conditions:

- Patients who cannot tolerate, including allergy or hypersensitivity to, procedural anticoagulation or post procedural anti-platelet regimen
- Patients with known hypersensitivity to clip components (nickel / titanium, cobalt, chromium, polyester), or with contrast sensitivity
- Active endocarditis of the mitral valve
- Rheumatic mitral valve disease
- Evidence of intracardiac, inferior vena cava (IVC) or femoral venous thrombus

### WARNINGS

- + DO NOT use MitraClip  $^{\scriptscriptstyle\rm TM}$  outside of the labeled indication.
- The MitraClip<sup>™</sup> G4 Implant should be implanted with sterile techniques using fluoroscopy and echocardiography (e.g., transesophageal [TEE] and transthoracic [TTE]) in a facility with on-site cardiac surgery and immediate access to a cardiac operating room.
- Read all instructions carefully. Use universal precautions for biohazards and sharps while handling the MitraClip<sup>™</sup> G4 System to avoid user injury. Failure to follow these instructions, warnings and precautions may lead to device damage, user injury or patient injury, including:
- MitraClip™ G4 Implant erosion, migration or malposition
- Failure to deliver MitraClip<sup>™</sup> G4 Implant to the intended site
  Difficulty or failure to retrieve MitraClip<sup>™</sup> G4 system components
- Use caution when treating patients with hemodynamic instability requiring inotropic support or mechanical heart assistance due to the increased risk of mortality in this patient population. The safety and effectiveness of MitraClip<sup>™</sup> in these patients has not been evaluated.
- Patients with a rotated heart due to prior cardiac surgery in whom the System is used may have a potential risk of experiencing adverse events such as atrial perforation, cardiac tamponade, tissue damage, and embolism which may be avoided with preoperative evaluation and proper device usage.
- For the Steerable Guide Catheter and Delivery Catheter only:
- The Guide Catheter: the distal 65 cm of the Steerable Guide Catheter with the exception of the distal soft tip, is coated with a hydrophilic coating.
- The Delivery Catheter: coated with a hydrophilic coating for a length of approximately 131 cm.
- Failure to prepare the device as stated in these instructions and failure to handle the device with care could lead to additional intervention or serious adverse event.
- The Clip Delivery System is provided sterile and designed for single use only. Cleaning, re-sterilization and / or re-use may result in infections, malfunction of the device and other serious injury or death.
- Note the product "Use by" date specified on the package.
- Inspect all product prior to use. Do not use if the package is open or damaged, or if product is damaged.

### PRECAUTIONS

- Prohibitive Risk Primary (or degenerative) Mitral Regurgitation
  - Prohibitive risk is determined by the clinical judgment of a heart team, including a cardiac surgeon experienced in mitral valve surgery and a cardiologist experienced in mitral valve disease, due to the presence of one or more of the following documented surgical risk factors:
  - 30-day STS predicted operative mortality risk score of
    - ${}_{\circ}\geq 8\%$  for patients deemed likely to undergo mitral valve replacement or
    - $\geq$  6% for patients deemed likely to undergo mitral valve repair
- Porcelain aorta or extensively calcified ascending aorta.
- Frailty (assessed by in-person cardiac surgeon consultation)
  Hostile chest
- Severe liver disease / cirrhosis (MELD Score > 12)
- Severe pulmonary hypertension (systolic pulmonary artery pressure > 2/3 systemic pressure)
- Unusual extenuating circumstance, such as right ventricular dysfunction with severe tricuspid regurgitation, chemotherapy for malignancy, major bleeding diathesis, immobility, AIDS, severe dementia, high risk of aspiration, internal mammary artery (IMA) at high risk of injury, etc.
- Evaluable data regarding safety or effectiveness is not available for prohibitive risk primary patients with an LVEF < 20% or an LVESD > 60 mm. MitraClip<sup>™</sup> should be used only when criteria for clip suitability for primary have been met.
- The heart team should include a cardiac surgeon experienced in mitral valve surgery and a cardiologist experienced in mitral valve disease and may also include appropriate physicians to assess the adequacy of heart failure treatment and valvular anatomy.
- · Secondary Mitral Regurgitation
  - Evaluable data regarding safety or effectiveness is not available for secondary MR patients with an LVEF < 20% or an LVESD > 70 mm.
- The multidisciplinary heart team should be experienced in the evaluation and treatment of heart failure and mitral valve disease and determine that symptoms and MR severity persist despite maximally tolerated GDMT.

### POTENTIAL COMPLICATIONS AND ADVERSE EVENTS

The following ANTICIPATED EVENTS have been identified as possible complications of the MitraClip<sup>™</sup> G4 procedure.

Death; Allergic reactions or hypersensitivity to latex, contrast agent, anaesthesia, device materials (nickel / titanium, cobalt, chromium, polyester), and drug reactions to anticoagulation, or antiplatelet drugs; Vascular access complications which may require transfusion or vessel repair including: wound dehiscence, catheter site reactions, Bleeding (including ecchymosis, oozing, hematoma, hemorrhage, retroperitoneal hemorrhage), Arteriovenous fistula, pseudoaneurysm, aneurysm, dissection, perforation / rupture, vascular occlusion, Emboli (air thrombotic material, implant, device component), Peripheral Nerve Injury; Lymphatic complications; Pericardial complications which may require additional intervention, including: Pericardial effusion, Cardiac tamponade, Pericarditis; Cardiac complications which may require additional interventions or emergency cardiac surgery, including: Cardiac perforation, Atrial septal defect; Mitral valve complications, which may complicate or prevent later surgical repair, including: Chordal entanglement / rupture, Single Leaflet Device Attachment (SLDA), Thrombosis, Dislodgement of previously implanted devices, Tissue damage, Mitral valve stenosis, Persistent or residual mitral regurgitation, Endocarditis; Cardiac arrhythmias (including conduction disorders, atrial arrhythmias, ventricular arrhythmias); Cardiac ischemic conditions (including myocardial infarction, myocardial ischemia, and unstable / stable angina); Venous thromboembolism (including deep vein thrombosis, pulmonary embolism, post procedure pulmonary embolism); Stroke / Cerebrovascular accident (CVA) and Transient Ischemic Attack (TIA); System organ failure: Cardio-respiratory arrest, Worsening heart failure, Pulmonary congestion, Respiratory dysfunction / failure / atelectasis, Renal insufficiency or failure, Shock (including cardiogenic and anaphylactic); Blood cell disorders (including coagulopathy, hemolysis, and Heparin Induced Thrombocytopenia (HIT)); Hypotension / hypertension; Infection including: Urinary Tract Infection (UTI), Pneumonia, Septicemia, Nausea / vomiting; Chest pain; Dyspnea; Edema; Fever or hyperthermia; Pain; Fluoroscopy, Transesophageal echocardiogram (TEE) and Transthoracic echocardiogram (TTE) -related complications: Skin injury or tissue changes due to exposure to ionizing radiation, Esophageal irritation, Esophageal perforation, Gastrointestinal bleeding.

CAUTION: These products are intended for use by or under the direction of a physician. Prior to use, reference the Instructions for Use, inside the product carton (when available) or at *eifu.abbottvascular.com* or at *medical.abbott/manuals* for more detailed information on Indications, Contraindications, Warnings, Precautions and Adverse Events.

Photos on file at Abbott.

Abbott

3200 Lakeside Dr., Santa Clara, CA 95054 USA, Tel: 1.800.227.9902

™ Indicates a trademark of the Abbott Group of Companies.

www.cardiovascular.abbott ©2020 Abbott. All rights reserved. MAT-2001330 v1.0 Item approved for U.S. use only.

